



CHEERLEADING INJURIES

Legend has it that cheerleading started with a University of Minnesota student standing up in the stands and leading his fellow students in “cheering” for their team during a football game. Cheerleading has morphed drastically since then. Today, it’s considered an athletic activity that incorporates elements of dance and gymnastics along with stunts and pyramid formations.

In 2002, an estimated 3.5 million people in the United States participated as cheerleaders, from six-year-olds to adults who cheerlead for professional athletic teams. While cheerleading is meant to support an athletic team, its intense competitions at the high school and collegiate levels have created a whole new dynamic, including increased risk for injury.

What types of injuries are most common in cheerleading?

The U.S. Consumer Product Safety Commission (CPSC) estimates that cheerleading led to 16,000 emergency room visits in 2002 (the latest year for data). While not as frequent as injuries in other sports, cheerleading injuries tend to be more severe, making up more than half of the catastrophic injuries in female athletes. Cheerleading injuries affect all areas of the body—most commonly the wrists, shoulders, ankles, head, and neck.

How can injuries be prevented?

Stunt Restrictions

In an attempt to curb the amount of catastrophic injuries in cheerleading, restrictions have been placed on stunts. They range from height restrictions in human pyramids, to the thrower-flyer ratio, to the number of spotters that must be present for each person lifted above shoulder level.

The limit for pyramids is two body lengths for the high school level and 2.5 body lengths for the college level, with the base cheerleader in direct contact with the performing surface. Base supporters must remain stationary and the suspended person is not allowed to be inverted or rotate on dismount. Basket toss stunts in which a cheerleader is thrown into the air (sometimes as high as 20 feet) are only allowed to have four throwers. The person being tossed (flyer) is not allowed to drop the head below a horizontal plane with the torso. One of the throwers must remain behind the flyer at all times during the toss.

CHEERLEADING INJURIES



Mats should be used during practice sessions and as much as possible during competitions. Cheerleaders should not attempt a stunt if they are tired, injured, or ill, as this may disrupt their focus and cause the stunt to be performed in an unsafe manner.

Training

The importance of a qualified coach is also critical. Coaching certification is encouraged.

Precautions should always be taken during inclement weather for all stunts. Also, a stunt should not be attempted without proper training, and not until the cheerleader is confident and comfortable with performing the stunt. Supervision should be provided at all times during stunt routines.

As with any sport, proper conditioning and training are important to minimize injury, including:

- Resistance exercises to gain strength in the lower back, stomach, and shoulders
- Regular stretching, yoga, or Pilates instruction to improve flexibility
- Speaking with a sports medicine professional or athletic trainer if you have any concerns about injuries or prevention strategies
- Returning to play only when clearance is granted by a healthcare professional

How are cheerleading injuries treated?

One of the most common injuries cheerleaders suffer is an ankle or wrist sprain or strain. For treatment of a sprain or strain, remember RICE:

- Rest the injured site for at least 24 hours
- Ice the injury at least every hour for 10–20 minutes during the initial four hours after injury. Icing then can be done 10–20 minutes four times a day for two days
- Compress the injured site with a snug, elastic bandage for 48 hours
- Elevate the injured limb for at least 24 hours X-rays may be required for more significant injuries.

Unfortunately, as cheerleading routines become more complex and acrobatic, cheerleaders are becoming more prone to significant head and neck injuries. Emergent medical attention with spine immobilization is required in these cases.

Expert Consultants

John D Campbell, MD
Barry P. Boden, MD

References

Campbell JD, Boden BP. Cheerleading injuries. *Sports Medicine Update*. September/October 2008.

Boden BP, Tacchetti R, Mueller FO. Catastrophic cheerleading injuries. *Am J Sports Med*. 31:881-888, 2003.

Shields BJ, Smith GA. Cheerleading-related injuries to children 5 to 18 years of age: United States, 1990-2002. *Pediatrics*. 117:122-129, 2006.

An AOSSM & STOP Sports Injuries Collaborating Organization



Sports Tips are brought to you by the American Orthopaedic Society for Sports Medicine. They provide general information only and are not a substitute for your own good judgment or consultation with a physician. To learn more about other orthopaedic sports medicine topics, visit sportsmed.org.

Copyright ©2022. American Orthopaedic Society for Sports Medicine. All rights reserved. Multiple copy reproduction prohibited without specific written permission.